

More on Penrose Limit of $AdS_4 \times Q^{1,1,1}$

Changhyun Ahn

*Department of Physics,
Kyungpook National University,
Taegu 702-701, Korea
ahn@knu.ac.kr*

abstract

We consider a Penrose limit of $AdS_4 \times Q^{1,1,1}$ that provides the pp-wave geometry equal to the one in the Penrose limit of $AdS_4 \times \mathbf{S}^7$. We expect that there exists a subsector of three dimensional $\mathcal{N} = 2$ dual gauge theory which has enhanced $\mathcal{N} = 8$ maximal supersymmetry. We identify operators in the $\mathcal{N} = 2$ gauge theory with 11-dimensional supergravity KK excitations in the pp-wave geometry and describe how both the chiral multiplets and semi-conserved multiplets fall into $\mathcal{N} = 8$ supermultiplets.

1 Introduction

Recently, it was found in [1] that the large N limit of a subsector of four dimensional $\mathcal{N} = 4$ $SU(N)$ supersymmetric gauge theory is dual to type IIB string theory in the pp-wave background [2, 3]. In the $N \rightarrow \infty$ limit with the finiteness of string coupling constant, this subspace of the gauge theory and the operator algebra are described by string theory in the pp-wave geometry. By considering a scale limit of the geometry near a null geodesic in $AdS_5 \times \mathbf{S}^5$, it leads to a truncation to the appropriate subspace of the gauge theory. The operators in the subsector of $\mathcal{N} = 4$ gauge theory can be identified with the excited string states in the pp-wave background. There are various and relevant papers [4]-[42] related to the work of [1].

There exist some results [7, 8, 10] on the Penrose limit of $AdS_5 \times T^{1,1}$ that gives the pp-wave geometry of $AdS_5 \times \mathbf{S}^5$. There is a subsector of $\mathcal{N} = 1$ gauge theory that contains an enhanced $\mathcal{N} = 4$ supersymmetry. The corresponding operators in the gauge theory side were identified with the stringy excitations in the pp-wave geometry and some of the gauge theory operators are combined into $\mathcal{N} = 4$ supersymmetry multiplets [8].

In this paper, we consider a similar duality that is present between a certain three dimensional $\mathcal{N} = 2$ gauge theory and 11-dimensional supergravity theory in a pp-wave background with the same spirit as in [7, 8, 10]. We describe this duality by taking a scaling limit of both 11-dimensional supergravity on $AdS_4 \times Q^{1,1,1}$ ($Q^{1,1,1}$ was found in [43] first to show an Einstein metric) and three dimensional superconformal field theory. Three-dimensional theory consists of an $\mathcal{N} = 2$ $SU(N)_1 \times SU(N)_2 \times SU(N)_3$ gauge theories with three kinds of chiral fields $A_i, i = 1, 2$ transforming in the $(\mathbf{N}, \overline{\mathbf{N}}, \mathbf{1})$ representation, $B_j, j = 1, 2$ transforming in the $(\mathbf{1}, \mathbf{N}, \overline{\mathbf{N}})$ representation and $C_k, k = 1, 2$ transforming in the $(\overline{\mathbf{N}}, \mathbf{1}, \mathbf{N})$ representation. This gives the theory of [45] that lives on N M2-branes at the conical singularity of a Calabi-Yau four-fold. The complete analysis on the spectrum of $AdS_4 \times Q^{1,1,1}$ was studied by Merlatti [44]. The scaling limit is obtained by considering the geometry near a null geodesic carrying large angular momentum in the $U(1)_R$ isometry of the $Q^{1,1,1}$ space which is dual to the $U(1)_R$ R-symmetry in the $\mathcal{N} = 2$ superconformal field theory. In section 2, we consider and review the scaling limit around a null geodesic in $AdS_4 \times Q^{1,1,1}$ and obtain an $\mathcal{N} = 8$ maximally supersymmetric pp-wave background of $AdS_4 \times \mathbf{S}^7$ described by [17]. In section 3, we identify supergravity KK excitations obtained by Merlatti [44] in the Penrose limit with gauge theory operators. In section 4, we summarize our results.

2 Penrose Limit of $AdS_4 \times Q^{1,1,1}$

Let us start with the supergravity solution dual to the $\mathcal{N} = 2$ superconformal field theory [45]. By putting a large number of N coincident M2-branes at the conifold singularity and taking

the near horizon limit, the metric becomes that [46, 45] of $AdS_4 \times Q^{1,1,1}$ (See also [47, 48, 49])

$$ds_{11}^2 = ds_{AdS_4}^2 + ds_{Q^{1,1,1}}^2, \quad (1)$$

where

$$\begin{aligned} ds_{AdS_4}^2 &= L^2 \left(-\cosh^2 \rho \, dt^2 + d\rho^2 + \sinh^2 \rho \, d\Omega_2^2 \right), \\ ds_{Q^{1,1,1}}^2 &= \frac{L^2}{16} \left(d\psi + \sum_{i=1}^3 \cos \theta_i d\phi_i \right)^2 + \frac{L^2}{8} \sum_{i=1}^3 \left(d\theta_i^2 + \sin^2 \theta_i d\phi_i^2 \right), \end{aligned}$$

where $d\Omega_2$ is the volume form of a unit two sphere $\mathbf{S}^2$ and the curvature radius¹ L of AdS_4 is given in terms of the number of M2-branes by $(2L)^6 = 32\pi^2 \ell_p^6 N$. Topologically $Q^{1,1,1}$ is a $U(1)$ bundle over $\mathbf{S}_1^2 \times \mathbf{S}_2^2 \times \mathbf{S}_3^2$. The base is parametrized by the spherical coordinates (θ_i, ϕ_i) , $i = 1, 2, 3$ to parametrize i -th two sphere, as usual, and the angle ψ which has period 4π parametrizes the $U(1)$ Hopf fiber. The $SU(2)_1 \times SU(2)_2 \times SU(2)_3 \times U(1)_R$ isometry group of $Q^{1,1,1}$ corresponds to $SU(2)_1 \times SU(2)_2 \times SU(2)_3$ global symmetry and $U(1)_R$ R -symmetry of the dual superconformal field theory of [45].

Let us make a scaling limit around a null geodesic in $AdS_4 \times Q^{1,1,1}$ that rotates along the ψ coordinate of $Q^{1,1,1}$ whose shift symmetry corresponds to the $U(1)_R$ symmetry in the dual superconformal field theory. Similar analysis in $AdS_5 \times T^{1,1}$ appeared in [7, 8, 10]. Let us introduce coordinates which label the geodesic

$$x^+ = \frac{1}{2} \left(t + \frac{1}{4} (\psi + \phi_1 + \phi_2 + \phi_3) \right), \quad x^- = \frac{L^2}{2} \left(t - \frac{1}{4} (\psi + \phi_1 + \phi_2 + \phi_3) \right),$$

and make a scaling limit around $\rho = 0 = \theta_1 = \theta_2 = \theta_3$ in the above geometry (1). By taking the limit $L \rightarrow \infty$ while rescaling the coordinates

$$\rho = \frac{r}{L}, \quad \theta_1 = \frac{\sqrt{2}\zeta_1}{L}, \quad \theta_2 = \frac{\sqrt{2}\zeta_2}{L}, \quad \theta_3 = \frac{\sqrt{2}\zeta_3}{L},$$

the Penrose limit of the $AdS_4 \times Q^{1,1,1}$ becomes [17]

$$\begin{aligned} ds_{11}^2 &= -4dx^+ dx^- + \sum_{i=1}^3 \left(dr^i dr^i - r^i r^i dx^+ dx^+ \right) + \frac{1}{4} \sum_{i=1}^3 \left(d\zeta_i^2 + \zeta_i^2 d\phi_i^2 - 2\zeta_i^2 d\phi_i dx^+ \right) \\ &= -4dx^+ dx^- + \sum_{i=1}^3 \left(dr^i dr^i - r^i r^i dx^+ dx^+ \right) + \frac{1}{4} \sum_{i=1}^3 \left(dz_i d\bar{z}_i + i(\bar{z}_i dz_i - z_i d\bar{z}_i) dx^+ \right) \quad (2) \end{aligned}$$

where in the last line we introduce the complex coordinates $z_i = \zeta_i e^{i\phi_i}$. Since the metric has a covariantly constant null Killing vector ∂/∂_{x^-} , it is also pp-wave metric. The pp-wave has a

¹Comparing with the one in [50], the L^6 here is related to $3L/8$ there. Note that the ratio of the volume of $\mathbf{S}^7$ to the volume of $Q^{1,1,1}$ is $8/3$. The overall factor $(8/3)^{1/3}$ is absorbed in the 11-dimensional metric ds_{11}^2 .

decomposition of the $\mathbf{R}^9$ transverse space into $\mathbf{R}^3 \times \mathbf{R}_1^2 \times \mathbf{R}_2^2 \times \mathbf{R}_3^2$ where $\mathbf{R}^3$ is parametrized by r^i and $\mathbf{R}_1^2 \times \mathbf{R}_2^2 \times \mathbf{R}_3^2$ by z_i . The symmetries of this background are the $SO(3)$ rotations in $\mathbf{R}^3$ and a $U(1)_1 \times U(1)_2 \times U(1)_3$ symmetry corresponding to the rotations of $\mathbf{R}^2 \times \mathbf{R}^2 \times \mathbf{R}^2$. In the gauge theory side, the $SO(3)$ symmetry corresponds to the subgroup of the $SO(2,3)$ conformal group and $U(1)_1 \times U(1)_2 \times U(1)_3$ rotation charges J_1, J_2 and J_3 correspond to the linear combination of Q_1 and R , Q_2 and R , and Q_3 and R respectively. R is the $U(1)_R$ charge of the gauge theory and Q_1, Q_2 and Q_3 are the Cartan generators of the $SU(2)_1 \times SU(2)_2 \times SU(2)_3$ global symmetry of the dual superconformal field theory. Note that the pp-wave geometry (2) in the scaling limit produces to the $\mathcal{N} = 8$ maximally supersymmetric pp-wave solution of $AdS_4 \times \mathbf{S}^7$ [51, 52, 18] through $z_i = e^{ix^+} w_i$ and $\bar{z}_i = e^{-ix^+} \bar{w}_i$ [8]

$$ds_{11}^2 = -4dx^+ dx^- + \sum_{i=1}^9 dr^i dr^i - \left(\sum_{i=1}^3 r^i r^i + \frac{1}{4} \sum_{i=4}^9 r^i r^i \right) dx^+ dx^+$$

after an $U(1)_1 \times U(1)_2 \times U(1)_3$ rotation in the $\mathbf{R}^2 \times \mathbf{R}^2 \times \mathbf{R}^2$ plane [17]. The supersymmetry enhancement in the Penrose limit implies that a hidden $\mathcal{N} = 8$ supersymmetry is present in the corresponding subsector of the dual $\mathcal{N} = 2$ superconformal field theory. In the next section, we provide a description of how to understand the excited KK states in the supergravity theory that corresponds in the dual superconformal field theory to operators with a given conformal dimension.

3 Gauge Theory Spectrum

The 11-dimensional supergravity theory in $AdS_4 \times Q^{1,1,1}$ is dual to the $\mathcal{N} = 2$ gauge theory encoded in a quiver diagram with gauge group $SU(N)_1 \times SU(N)_2 \times SU(N)_3$ with three series of chiral fields $A_i, i = 1, 2$ transforming in the $(\mathbf{N}, \overline{\mathbf{N}}, \mathbf{1})$ color representation, $B_j, j = 1, 2$ transforming in the $(\mathbf{1}, \mathbf{N}, \overline{\mathbf{N}})$ color representation and $C_k, k = 1, 2$ transforming in the $(\overline{\mathbf{N}}, \mathbf{1}, \mathbf{N})$ color representation [45]. At the fixed point, these chiral superfields have conformal weight $1/3$ and transform as $(\mathbf{2}, \mathbf{1}, \mathbf{1}), (\mathbf{1}, \mathbf{2}, \mathbf{1})$ and $(\mathbf{1}, \mathbf{1}, \mathbf{2})$ under the $SU(2)_1 \times SU(2)_2 \times SU(2)_3$ global symmetry. Chiral fields A_1 and A_2 are doublets of $SU(2)_1$ global symmetry, B_1 and B_2 are doublets of $SU(2)_2$ global symmetry, and C_1 and C_2 are doublets of $SU(2)_3$ global symmetry. There exist three kinds of supermultiplets in an $\mathcal{N} = 2$ AdS_4 superspace: massless multiplets, short multiplets and long multiplets [53]. We identify states in the supergravity with operators in the gauge theory and focus only on the bosonic excitations of the theory. In each multiplet, we specify field contents, a representation with $U(1)$ charge, conformal weight and $U(1)_R$ R -charge.

- **Massless multiplets** [53, 44]

1) Massless graviton multiplet: $^2 (2, \frac{3}{2}, \frac{3}{2}, 1), \quad (\mathbf{1}, \mathbf{1}, \mathbf{1})_0, \quad \Delta = 2, \quad R = 0.$

There exists a stress-energy tensor superfield $T_{\alpha\beta}(x, \theta)$ that satisfies the equation for conserved current $D_\alpha^+ T^{\alpha\beta}(x, \theta) = D_\alpha^- T^{\alpha\beta}(x, \theta) = 0$.³ This $T_{\alpha\beta}(x, \theta)$ is a singlet with respect to the flavor group $SU(2)_1 \times SU(2)_2 \times SU(2)_3$ and its conformal dimension is 2. Moreover R charge is 0. So this corresponds to the massless graviton multiplet that propagates in the AdS_4 bulk [45].

2) Massless vector multiplet: $(1, 1, 1, 0, 0), \quad (\mathbf{3}, \mathbf{1}, \mathbf{1})_0, \quad \Delta = 1, \quad R = 0.$

There exists a conserved vector current, a scalar superfield $J_{SU(2)_i}(x, \theta)$, to each generator of the flavor symmetry group through Noether theorem satisfying the conservation equations $D^{+\alpha} D_\alpha^+ J_{SU(2)_i}(x, \theta) = D^{-\alpha} D_\alpha^- J_{SU(2)_i}(x, \theta) = 0$. This $J_{SU(2)_1}(x, \theta)$ transforms in the adjoint representation of the first factor $SU(2)_1$ of the flavor group and its conformal dimension is 1 with vanishing R -charge. This corresponds to the massless vector multiplet propagating in the AdS_4 bulk [45]. There are also two massless vector multiplets denoted by $(\mathbf{1}, \mathbf{3}, \mathbf{1})_0$ and $(\mathbf{1}, \mathbf{1}, \mathbf{3})_0$ coming from the permutation of three $SU(2)$ flavor groups.

3) Two massless vector multiplets: $(1, 1, 1, 0, 0), \quad (\mathbf{1}, \mathbf{1}, \mathbf{1})_0, \quad \Delta = 1, \quad R = 0.$

It is known that two Betti currents of $Q^{1,1,1}$ are written down from the toric description [45] and they are conserved. Their conformal dimension is 1 and R -charge is zero. Therefore Betti currents correspond to additional massless vector multiplets. Therefore massless multiplets 1), 2) and 3) saturate the unitary bound and have a conformal weight(or dimension) related to the maximal spin.

• Short multiplets [53, 44]

It is known that the dimension of the scalar operator in terms of energy labels, in the dual SCFT corresponding $AdS_4 \times Q^{1,1,1}$ is

$$\Delta = \frac{3}{2} + \frac{1}{2} \sqrt{1 + \frac{m^2}{4}} = \frac{3}{2} + \frac{1}{2} \sqrt{45 + \frac{E}{4} - 6\sqrt{36 + E}}. \quad (3)$$

The first equation in (3) comes from the relation between the lowest energy eigenvalue and the mass which appears in the AdS_4 wave equation. The relation of d'Alembertian in AdS_4 to Casimir operator was obtained in [54, 55]. The second relation in (3) comes from the formula of mass $m^2 = E + 176 - 24\sqrt{E + 36}$ in [56] where the normalization for this is $(\square - 32 + m^2)S = 0$ for scalar field S . The energy spectrum on $Q^{1,1,1}$ exhibits an interesting feature which is relevant to superconformal algebra and it is given by [57, 50]

$$E = 32 \left(\sum_{i=1}^3 l_i(l_i + 1) - s^2 \right) \quad (4)$$

²Here we denote the field contents of each multiplet by their spin [46]. For massless graviton multiplet $(2, \frac{3}{2}, \frac{3}{2}, 1)$, there are one spin 2, two spin 3/2 and one spin 1.

³For three dimensional $\mathcal{N} = 2$ superfields, we define $\theta_\alpha^\pm = \frac{1}{\sqrt{2}} (\theta_\alpha^1 \pm i\theta_\alpha^2)$ and the superderivatives are defined as $D^{\pm\alpha} = \frac{\partial}{\partial \theta_\alpha^\pm} + \frac{1}{2} (\gamma^m)_\beta^\alpha \theta^\beta \partial_m$.

where the eigenvalue E is classified by $U(1)$ charge s and spin(or angular momentum) l_i 's under $SU(2)_1 \times SU(2)_2 \times SU(2)_3$ and the eigenmodes occur in $(\mathbf{2l_1} + \mathbf{1}, \mathbf{2l_2} + \mathbf{1}, \mathbf{2l_3} + \mathbf{1})_s$ dimensional representation and $l_i \geq |s|$, $s = 0, \pm\frac{1}{2}, \pm 1, \dots$. The $U(1)$ part of the isometry group of $Q^{1,1,1}$ acts by shifting $U(1)$ charge s . The integer R -charge, R is related to $U(1)$ charge s by $s = R/2$. Let us take $R \geq 0$. One can find the lowest value of Δ is R corresponding to a mode scalar with $l_i = s$ because E becomes $32(2s^2 + 3s)$ and by plugging back to (3) then one obtains $\Delta = R$. Thus we find a set of operators filling out a $(\mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{1})_{\frac{R}{2}}$ multiplet of $SU(2)_1 \times SU(2)_2 \times SU(2)_3 \times U(1)$ where a subscript is $U(1)$ charge $R/2$ and the number $\mathbf{R} + \mathbf{1}$ is the dimension of each $SU(2)$ representation. The condition $\Delta = R$ saturates the bound on Δ from superconformal algebra. The fact that the R -charge of a chiral operator is equal to the conformal dimension was observed in [6] in the context of R symmetry gauge field.

1) One hypermultiplet: $(\frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0)$

$$(\mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{1})_{\frac{R}{2}}, \quad \Delta = R.$$

It was shown in [45] that from the harmonic analysis on $Q^{1,1,1}$ and the spectrum of $SU(2)_1 \times SU(2)_2 \times SU(2)_3$ representation of the $OSp(2|4)$ hypermultiplets, the hypermultiplet of conformal dimension $\Delta = R$ and $U(1)$ charge $s = R/2$ should be in the representation $l_i = s = R/2$. According to [53], the information on the Laplacian eigenvalues allows us to get the spectrum of hypermultiplets of the theory corresponding to the chiral operators of the SCFT. This part of spectrum was given in [45] and the form of operators is

$$\text{Tr}\Phi_C \equiv \text{Tr}(ABC)^R \tag{5}$$

where the flavor $SU(2)$'s indices are totally symmetrized and the chiral superfield $\Phi_C(x, \theta)$ satisfies $D_\alpha^+ \Phi_C(x, \theta) = 0$. The hypermultiplet spectrum in the KK harmonic expansions on $Q^{1,1,1}$ agrees with the chiral superfield predicted by the conformal gauge theory. Based on the work of [55] on a generic $OSp(2|4)$ representation, the operators with protected dimensions are related to shortenings and they are classified as three categories: 1) chiral superfields, 2) conserved currents and 3) semi-conserved currents. The definitions of these fields are given in [58]. From this, the dimension of ABC should be 1 to match the spectrum. One realizes that the chiral fields A_i, B_j, C_k have conformal weight $1/3$.

2) One short graviton multiplet: $(2, \frac{3}{2}, \frac{3}{2}, \frac{3}{2}, 1, 1, 1, \frac{1}{2})$ [44]

$$(\mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{1})_{\frac{R}{2}}, \quad \Delta = R + 2.$$

The gauge theory interpretation of this multiplet is obtained by adding a dimension 2 singlet operator with respect to flavor group into the above chiral superfield $\Phi_C(x, \theta)$. We consider $\text{Tr}\Phi_{\alpha\beta} \equiv \text{Tr}(T_{\alpha\beta}\Phi_C)$, where $T_{\alpha\beta}(x, \theta)$ is a stress energy tensor we have discussed

before and $\Phi_{\text{C}}(x, \theta)$ is a chiral superfield in (5). All color indices are symmetrized before taking the contraction. This composite operator satisfies $D_{\alpha}^{+} \Phi^{\alpha\beta}(x, \theta) = 0$.

3) One short vector multiplet: $(1, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0)$ [44]

$$(\mathbf{R} + \mathbf{3}, \mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{1})_{\frac{\mathbf{R}}{2}}, \quad \Delta = R + 1.$$

One can construct the following gauge theory object corresponding to the short vector multiplet: $\text{Tr} \Phi \equiv \text{Tr} (J_{SU(2)_1} \Phi_{\text{C}})$, where $J_{SU(2)_1}(x, \theta)$ is a conserved vector current transforming in the adjoint representation of $SU(2)_1$ flavor group as before in massless vector multiplet and $\Phi_{\text{C}}(x, \theta)$ is a chiral superfield in (5). There exist two other short multiplets by $(\mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{3}, \mathbf{R} + \mathbf{1})_{\frac{\mathbf{R}}{2}}$ and $(\mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{3})_{\frac{\mathbf{R}}{2}}$ due to the permutation of flavor groups. In this case, we have $D^{+\alpha} D_{\alpha}^{+} \Phi(x, \theta) = 0$. There is another short vector multiplet described by $(\mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{1})_{\frac{\mathbf{R}}{2}}, \Delta = R$. There exist three short gravitino multiplets $(\frac{3}{2}, 1, 1, 1, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0)$ [44] specified by

$$(\mathbf{R}, \mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{1})_{\frac{\mathbf{R}}{2}+1}, \quad (\mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{2}, \mathbf{R} + \mathbf{2})_{\frac{\mathbf{R}}{2}-1}, \quad (\mathbf{R}, \mathbf{R} + \mathbf{1}, \mathbf{R} + \mathbf{1})_{\frac{\mathbf{R}}{2}+1}$$

with conformal weight, $\Delta = R + 3/2$. Therefore the short multiplets 1), 2) and 3) saturate the unitary bound and have a conformal dimension related to the R -charge and maximal spin. In superfield language, the θ expansion of the superfield is shortened by imposing a suitable differential constraint, invariant with respect to Poincare supersymmetry.

• Long multiplets [44]

Although the dimensions of nonchiral operators are in general irrational, there exist special integer values of k_i such that for $l_i = k_i + R/2$, one can see the Diophantine like condition [50],

$$-2(k_1 k_2 + k_2 k_3 + k_3 k_1) + \sum_{i=1}^3 (k_i^2 - k_i) = 0 \quad (6)$$

make $\sqrt{36 + E}$ be equal to $4R + 2(2\sum_{i=1}^3 k_i + 3)$. This resembles the one in $AdS_5 \times T^{1,1}$ [8]. Furthermore in order to make the dimension be rational (their conformal dimensions are protected), $45 + E/4 - 6\sqrt{36 + E}$ in (3) should be square of something. It turns out this is the case without any further restrictions on k_i 's. Therefore we have $\Delta = R + \sum_i^3 k_i$ which is Δ_+ for $\Delta \geq 3/2$ and Δ_- for $\Delta \leq 3/2$. This is true if we are describing states with finite Δ and R . Since we are studying the scaling limit $\Delta, R \rightarrow \infty$, we have to modify the above analysis. This constraint (6) comes from the fact that the energy eigenvalue of the Laplacian on $Q^{1,1,1}$ for the supergravity mode (4) takes the form

$$E = \sum_{i=1}^3 (k_i^2 + (R + 1)k_i) + \frac{R}{2} (R + 3). \quad (7)$$

One can show that the conformal weight of the long vector multiplet A below becomes rational if the condition (6) is satisfied. This multiplet has the following properties.

1) One long vector multiplet A : $(1, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, 0)$

$$(2\mathbf{k}_1 + \mathbf{R} + \mathbf{1}, 2\mathbf{k}_2 + \mathbf{R} + \mathbf{1}, 2\mathbf{k}_3 + \mathbf{R} + \mathbf{1})_{\frac{\mathbf{R}}{2}}, \quad \Delta = -\frac{3}{2} + \frac{1}{4}\sqrt{E + 36}.$$

However, as we take the limit of $R \rightarrow \infty$, this constraint (6) is relaxed. The combination of $\Delta - R$ is given by

$$\Delta - R = k_1 + k_2 + k_3 + \mathcal{O}\left(\frac{1}{R}\right) \quad (8)$$

where the right hand side is definitely rational and they are integers. So the constraint (6) is not relevant in the subsector of the Hilbert space we are interested in. Candidates for such states in the gauge theory side are given in terms of semi-conserved superfields [45]. Although they are not chiral primaries, their conformal dimensions are protected. The ones we are interested in take the following form,

$$\text{Tr}\Phi_{\text{s.c.}} \equiv \text{Tr} \left[\left(J_{SU(2)_1} \right)^{k_1} \left(J_{SU(2)_2} \right)^{k_2} \left(J_{SU(2)_3} \right)^{k_3} (ABC)^R \right] \quad (9)$$

where the scalar superfields $J_{SU(2)_i}(x, \theta)$ transform in the adjoint representation of flavor group $SU(2)_i$ and satisfy $D^{+\alpha} D_{\alpha}^{+} J_{SU(2)_i}(x, \theta) = D^{-\alpha} D_{\alpha}^{-} J_{SU(2)_i}(x, \theta) = 0$ with conformal dimension 1 and zero $U(1)_R$ charge. Also we have $D^{+\alpha} D_{\alpha}^{+} \Phi_{\text{s.c.}}(x, \theta) = 0$. Since the singleton superfields $A_{i,b}^a$ carry an index a in the $\mathbf{N}$ of $SU(N)_1$ and an index b in the $\overline{\mathbf{N}}$ of the $SU(N)_2$, the fields $B_{j,c}^b$ carry an index b in the $\mathbf{N}$ of $SU(N)_2$ and an index c in the $\overline{\mathbf{N}}$ of the $SU(N)_3$, and $C_{k,a}^c$ carry an index c in the $\mathbf{N}$ of $SU(N)_3$ and an index a in the $\overline{\mathbf{N}}$ of the $SU(N)_1$, one can construct the following conserved flavor currents transforming $(\mathbf{3}, \mathbf{1}, \mathbf{1})_0$, $(\mathbf{1}, \mathbf{3}, \mathbf{1})_0$ and $(\mathbf{1}, \mathbf{1}, \mathbf{3})_0$ respectively

$$\begin{aligned} \left(J_{SU(2)_1} \right)_{i_1}^{j_1} &= A^{j_1} \overline{A}_{i_1} - \frac{\delta_{i_1}^{j_1}}{2} A \overline{A}, \\ \left(J_{SU(2)_2} \right)_{i_2}^{j_2} &= B^{j_2} \overline{B}_{i_2} - \frac{\delta_{i_2}^{j_2}}{2} B \overline{B}, \\ \left(J_{SU(2)_3} \right)_{i_3}^{j_3} &= C^{j_3} \overline{C}_{i_3} - \frac{\delta_{i_3}^{j_3}}{2} C \overline{C}, \end{aligned}$$

where the color indices are contracted in the right hand side. Note that the conformal dimension of these currents is not the one of naive sum of A and $\overline{A}$, B and $\overline{B}$ and C and $\overline{C}$. As we discussed in the last section, supergravity theory in $AdS_4 \times Q^{1,1,1}$ acquires an enhanced $\mathcal{N} = 8$ superconformal symmetry in the Penrose limit. This implies that the spectrum of the gauge theory operators in this subsector should fall into $\mathcal{N} = 8$ multiplets. We expect that both the chiral primary fields of the form $\text{Tr}(ABC)^R$ and the semi-conserved multiplets of the form (9) combine into make $\mathcal{N} = 8$ multiplets in this limit. Note that for finite R , the semi-conserved multiplets should obey the Diophantine constraint (6) in order for them to possess rational conformal weights.

In the remaining multiplets we consider the following particular representations in the global symmetry group with $U(1)$ charge: $(2\mathbf{k}_1 + \mathbf{R} + 1, 2\mathbf{k}_2 + \mathbf{R} + 1, 2\mathbf{k}_3 + \mathbf{R} + 1)_{\frac{\mathbf{R}}{2}}$,

2) One long vector multiplet Z : $(1, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, 0)$, $\Delta = \frac{1}{2} + \frac{1}{4}\sqrt{E + 32R + 4}$

For finite R , in order to make the dimension of this multiplet rational the expression $E + 32R + 4$ should be square of something. By plugging the expression of E (7), it turns out $\sqrt{E + 32R + 4} = 4R + 4(k_1 + k_2 + k_3) + 2$ with same constraint (6). So the combination of $\Delta - R$ becomes

$$\Delta - R = 1 + k_1 + k_2 + k_3 + \mathcal{O}\left(\frac{1}{R}\right).$$

Since we do not have any singlet of conformal dimension 1 with respect to the flavor group, one cannot increase a conformal dimension by simply tensoring any extra superfields into a semi-conserved current in order to match the spectrum. So the only way to do this is to increase the number of conserved scalar superfield. In order to produce the following gauge theory operator $\text{Tr}\left(J_{SU(2)_1}\Phi_{\text{s.c.}}\right)$ corresponding to this vector multiplet, one can think of a higher dimensional representation in the first factor $SU(2)_1$ in the global symmetry. Then the constraint coming from the requirement of rationality of conformal dimension is also changed for finite Δ and R .

3) One long graviton multiplet h : $(2, \frac{3}{2}, \frac{3}{2}, \frac{3}{2}, \frac{3}{2}, 1, 1, 1, 1, 1, 1, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0)$

$$\Delta = \frac{1}{2} + \frac{1}{4}\sqrt{E + 36}.$$

For finite R with rational dimension, after inserting the E into the above, we will arrive at the relation with same constraint (6) which is greater than (8) by 2:

$$\Delta - R = 2 + k_1 + k_2 + k_3 + \mathcal{O}\left(\frac{1}{R}\right). \quad (10)$$

The gauge theory interpretation of this multiplet is quite simple. If we take a semi-conserved current $\Phi_{\text{s.c.}}(x, \theta)$ defined in (9) and multiply it by a stress-energy tensor superfield $T_{\alpha\beta}(x, \theta)$ that is a singlet with respect to the flavor group, namely

$$\text{Tr}(T_{\alpha\beta}\Phi_{\text{s.c.}}),$$

we reproduce the right $OSp(2|4) \times SU(2)_1 \times SU(2)_2 \times SU(2)_3$ representations of the long vector multiplet. Also one can expect that other candidate for this multiplet with different representation by multiplying a semi-conserved current into a quadratic conserved scalar superfield: $\text{Tr}\left(J_{SU(2)_i}J_{SU(2)_j}\Phi_{\text{s.c.}}\right)$. The constraint for finite Δ and R is shifted as $k_i \rightarrow k_i + 1$ and $k_j \rightarrow k_j + 1$.

4) One long vector multiplet W : $(1, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, 0)$, $\Delta = \frac{5}{2} + \frac{1}{4}\sqrt{E + 36}$.

In this case, we get $\Delta - R$ by adding 2 to the one in (10)

$$\Delta - R = 4 + k_1 + k_2 + k_3 + \mathcal{O}\left(\frac{1}{R}\right).$$

One can describe corresponding gauge theory operator by taking quadratic stress-energy tensor $T_{\alpha\beta}T^{\alpha\beta}(x, \theta)$ and multiplying it into a semi-conserved current $\Phi_{\text{s.c.}}(x, \theta)$ we have seen before in order to match with the conformal dimension. That is, one obtains

$$\text{Tr} \left(T_{\alpha\beta} T^{\alpha\beta} \Phi_{\text{s.c.}} \right).$$

Similarly one can construct the following gauge theory operators related to this vector multiplet $\text{Tr} \left(T_{\alpha\beta} J_{SU(2)_i} J_{SU(2)_j} \Phi_{\text{s.c.}} \right)$ or $\text{Tr} \left(J_{SU(2)_i} J_{SU(2)_j} J_{SU(2)_k} J_{SU(2)_l} \Phi_{\text{s.c.}} \right)$. In the former, one can see the shift of $k_{i,j} \rightarrow k_{i,j} + 1$ and for the latter, the shift of $k_{i,j,k,l} \rightarrow k_{i,j,k,l} + 1$.

5) One long vector multiplet Z : $(1, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, 0)$, $\Delta = \frac{1}{2} + \frac{1}{4}\sqrt{E+4}$

Although there exists no rational dimension for this case with any choice of k_i 's when Δ and R are finite, the combination of $\Delta - R$ with Penrose limit $R \rightarrow \infty$ in the gauge theory side becomes

$$\Delta - R = 2 + k_1 + k_2 + k_3 + \mathcal{O}\left(\frac{1}{R}\right).$$

In addition to the above 1), 2), 3), 4) and 5) multiplets, there are also two long gravitino multiplets [44] $\chi^+(\frac{3}{2}, 1, 1, 1, 1, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0)$ characterized by $\Delta = -\frac{1}{2} + \frac{1}{4}\sqrt{E \pm 16R + 32}$ and two long gravitino multiplets χ^- characterized by $\Delta = \frac{3}{2} + \frac{1}{4}\sqrt{E \pm 16R + 32}$. Similar analysis can be done in this case. Although for finite Δ and R , both do not provide rational conformal dimensions, in the Penrose limit there is no constraint on the integer values and $R \rightarrow \infty$ limit will give us a rational conformal dimension.

4 Conclusion

We described an explicit example of an $\mathcal{N} = 2$ three-dimensional superconformal field theory that has a subsector of the Hilbert space with enhanced $\mathcal{N} = 8$ superconformal symmetry, in the large N limit. This subsector of gauge theory is achieved by Penrose limit which constrains strictly the states of the gauge theory to those whose conformal dimension and R charge diverge in the large N limit but possess finite value $\Delta - R$. We predicted for the spectrum of $\Delta - R$ of the $\mathcal{N} = 2$ superconformal field theory and proposed how the excited states in the supergravity correspond to gauge theory operators. In particular, both the chiral multiplets (5) and semi-conserved multiplets (9) of $\mathcal{N} = 2$ supersymmetry should combine into $\mathcal{N} = 8$ chiral multiplets. It would be interesting to study other types of compactification on M-theory in the Penrose limit which will provide pp-wave background and see what the corresponding gauge theory operators possessing a hidden $\mathcal{N} = 8$ supersymmetry are.

Acknowledgements

This research was supported by grant 2000-1-11200-001-3 from the Basic Research Program of the Korea Science & Engineering Foundation.

References

- [1] D. Berenstein, J. Maldacena and H. Nastase, JHEP **0204** (2002) 013, [hep-th/0202021](#).
- [2] M. Blau, J. Figueroa-O’Farrill, C.M. Hull and G. Papadopoulos, [hep-th/0201081](#).
- [3] M. Blau, J. Figueroa-O’Farrill, C.M. Hull and G. Papadopoulos, JHEP **0201** (2001) 047, [hep-th/0110242](#).
- [4] R.R. Metsaev and A.A. Tseytlin, [hep-th/0202109](#).
- [5] M. Blau, J. Figueroa-O’Farrill and G. Papadopoulos, [hep-th/0202111](#).
- [6] D. Berenstein, C.P. Herzog and I.R. Klebanov, [hep-th/0202150](#).
- [7] N. Itzhaki, I.R. Klebanov and S. Mukhi, JHEP **0203** (2002) 048, [hep-th/0202153](#).
- [8] J. Gomis and H. Ooguri, [hep-th/0202157](#).
- [9] J.G. Russo and A.A. Tseytlin, JHEP **0204** (2002) 021, [hep-th/0202179](#).
- [10] L.A. Pando Zayas and J. Sonnenschein, [hep-th/0202186](#).
- [11] M. Hatsuda, K. Kamimura and M. Sakaguchi, [hep-th/0202190](#).
- [12] M. Alishahiha and M.M. Sheikh-Jabbari, [hep-th/0203018](#).
- [13] M. Billo and I. Pesando, [hep-th/0203028](#).
- [14] N. Kim, A. Pankiewicz, S.-J. Rey and S. Theisen, [hep-th/0203080](#).
- [15] M. Cvetič, H. Lu and C.N. Pope, [hep-th/0203082](#).
- [16] T. Takayanagi and S. Terashima, [hep-th/0203093](#).
- [17] U. Gursoy, C. Nunez and M. Schwelling, [hep-th/0203124](#).
- [18] E. Floratos and A. Kehagias, [hep-th/0203134](#).
- [19] J. Michelson, [hep-th/0203140](#).
- [20] R. Gueven, [hep-th/0203153](#).

- [21] S.R. Das, C. Gomez and S.-J. Rey, [hep-th/0203164](#).
- [22] C.-S. Chu and P.-M. Ho, [hep-th/0203186](#).
- [23] M. Cvetič, H. Lu and C.N. Pope, [hep-th/0203229](#).
- [24] A. Dabholkar and S. Parvizi, [hep-th/0203231](#).
- [25] D. Berenstein, E. Gava, J. Maldacena, K.S. Narain and H. Nastase, [hep-th/0203249](#).
- [26] J.P. Gauntlett and C.M. Hull, [hep-th/0203255](#).
- [27] P. Lee and J. Park, [hep-th/0203257](#).
- [28] H. Lu and J.F. Vazquez-Poritz, [hep-th/0204001](#).
- [29] M. Hatsuda, K. Kamimura and M. Sakaguchi, [hep-th/0204002](#).
- [30] E. Kiritsis and B. Pioline, [hep-th/0204004](#).
- [31] A. Kumar, R.R. Nayak and Sanjay, [hep-th/0204025](#).
- [32] R.G. Leigh, K. Okuyama and M. Rozali, [hep-th/0204026](#).
- [33] D. Bak, [hep-th/0204033](#).
- [34] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, [hep-th/0204051](#).
- [35] K. Skenderis and M. Taylor, [hep-th/0204054](#).
- [36] M. Spradlin and A. Volovich, [hep-th/0204146](#).
- [37] S. Mukhi, M. Rangamani and E. Verlinde, [hep-th/0204147](#).
- [38] M. Alishahiha and M.M. Sheikh-Jabbari, [hep-th/0204174](#).
- [39] V. Balasubramanian, M. Huang, T.S. Levi and A. Naqvi, [hep-th/0204196](#).
- [40] Y. Imamura, [hep-th/0204200](#).
- [41] S. Frolov and A.A. Tseytlin, [hep-th/0204226](#).
- [42] H. Takayanagi and T. Takayanagi, [hep-th/0204234](#).
- [43] R. D'Auria, P. Fre and P. van Nieuwenhuizen, *Phys.Lett.* **B136** (1984) 347.
- [44] P. Merlatti, *Class.Quant.Grav.* **18** (2001) 2797, [hep-th/0012159](#).

- [45] D. Fabbri, P. Fre, L. Gualtieri, C. Reina, A. Tomasiello, A. Zaffaroni and A. Zampa, Nucl.Phys. **B577** (2000) 547, [hep-th/9907219](#).
- [46] D. Fabbri, P. Fre, L. Gualtieri and P. Termonia, Class.Quant.Grav. **17** (2000) 55, [hep-th/9905134](#).
- [47] B.E.W. Nilsson and C.N. Pope, Class.Quant.Grav. **1** (1984) 499.
- [48] D. Page and C.N. Pope, Phys.Lett. **B144** (1984) 346.
- [49] D. Page and C.N. Pope, Phys.Lett. **B145** (1984) 337.
- [50] C. Ahn, Phys.Lett. **B466** (1999) 171, [hep-th/9908162](#).
- [51] J. Kowalski-Glikman, Phys.Lett. **B134** (1984) 194.
- [52] J. Figueroa-O’Farrill and G. Papadopoulos, JHEP **0108** (2001) 036, [hep-th/0105308](#).
- [53] D. Fabbri, P. Fre, L. Gualtieri and P. Termonia, Nucl.Phys. **B560** (1999) 617, [hep-th/9903036](#).
- [54] H. Nicolai, Representations of Supersymmetry in Anti-De Sitter Space, Trieste School 1984:368 (QC178:S81:1984).
- [55] A. Ceresole, P. Fre and H. Nicolai, Class.Quant.Grav. **2** (1985) 133.
- [56] R. D’Auria and P. Fre, Annals Phys. **162** (1985) 372.
- [57] C.N. Pope, Class.Quant.Grav. **1** (1984) L91.
- [58] A. Ceresole, G. Dall’Agata, R. D’Auria and S. Ferrara, JHEP **0003** (2000) 011, [hep-th/9912107](#).